

LAW of COSINES DERIVATION

Sides AD and AP are tangent at A such that...

- DAP = spherical angle A
- DAO = 90°
- PAO = 90°

In plane triangle DPO...

Equation 1  $DP^2 = PO^2 + DO^2 - 2 \cdot PO \cdot DO \cdot \cos \alpha$

In plane triangle ADP...

Equation 2  $DP^2 = AD^2 + AP^2 - 2 \cdot AD \cdot AP \cdot \cos A$

In right triangle APO...

Equation 3  $AO^2 = PO^2 - AP^2$

In right triangle ADO...

Equation 4  $AO^2 = DO^2 - AD^2$

Equate the right sides of equations 1 and 2 and rearrange...

$$PO^2 + DO^2 - 2 \cdot PO \cdot DO \cdot \cos \alpha = AD^2 + AP^2 - 2 \cdot AD \cdot AP \cdot \cos A$$

Equation 5  $PO^2 + DO^2 - AD^2 - AP^2 = 2 \cdot PO \cdot DO \cdot \cos \alpha - 2 \cdot AD \cdot AP \cdot \cos A$

Add equations 3 and 4 and rearrange...

$$AO^2 + AO^2 = PO^2 - AP^2 + DO^2 - AD^2$$

Equation 6  $2 \cdot AO^2 = PO^2 + DO^2 - AD^2 - AP^2$


Equate the left side of equation 6 to the right side of equation 5 and rearrange...

$$2 \cdot AO^2 = 2 \cdot PO \cdot DO \cdot \cos \alpha - 2 \cdot AD \cdot AP \cdot \cos A$$

$$AO^2 + AD \cdot AP \cdot \cos A = PO \cdot DO \cdot \cos \alpha$$

$$\cos \alpha = \frac{AO \cdot AO}{PO \cdot DO} + \frac{AP \cdot AD}{PO \cdot DO} \cos A = \cos \mu \cdot \cos \beta + \sin \mu \cdot \sin \beta \cdot \cos A$$

note...  $\alpha = a, \beta = b, \mu = c$



Law of Cosines...

$$\cos a = \cos b \cdot \cos c + \sin b \cdot \sin c \cdot \cos A$$

$$\cos b = \cos a \cdot \cos c + \sin a \cdot \sin c \cdot \cos B$$

$$\cos c = \cos a \cdot \cos b + \sin a \cdot \sin b \cdot \cos C$$

} alternate versions

LAW of SINES DERIVATION

<u>Equation 7</u>	$\cos a = \cos b \cdot \cos c + \sin b \cdot \sin c \cdot \cos A$	}	by Law of Cosines
<u>Equation 8</u>	$\cos b = \cos a \cdot \cos c + \sin a \cdot \sin c \cdot \cos B$		
<u>Equation 9</u>	$\cos c = \cos a \cdot \cos b + \sin a \cdot \sin b \cdot \cos C$		

Rearrange equations 7, 8 and 9...

<u>Equation 10</u>	$\cos a - \cos b \cdot \cos c = \cos A \cdot \sin b \cdot \sin c$	( from 7 )
<u>Equation 11</u>	$\cos b - \cos a \cdot \cos c = \cos B \cdot \sin a \cdot \sin c$	( from 8 )
<u>Equation 12</u>	$\cos c - \cos a \cdot \cos b = \cos C \cdot \sin a \cdot \sin b$	( from 9 )

Using equation 10, square both sides...

$$\cos^2 a - 2 \cdot \cos a \cdot \cos b \cdot \cos c + \cos^2 b \cdot \cos^2 c = \cos^2 A \cdot \sin^2 b \cdot \sin^2 c$$

Apply the trig identity  $\cos^2 \theta = 1 - \sin^2 \theta \dots$

$$(1 - \sin^2 a) - 2 \cdot \cos a \cdot \cos b \cdot \cos c + (1 - \sin^2 b) \cdot (1 - \sin^2 c) = (1 - \sin^2 A) \cdot \sin^2 b \cdot \sin^2 c$$

$$1 - \sin^2 a - 2 \cdot \cos a \cdot \cos b \cdot \cos c + 1 - \sin^2 c - \sin^2 b + \sin^2 b \cdot \sin^2 c = \sin^2 b \cdot \sin^2 c - \sin^2 A \cdot \sin^2 b \cdot \sin^2 c$$

$$\text{Equation 13} \quad 2 - \sin^2 a - \sin^2 b - \sin^2 c - 2 \cdot \cos a \cdot \cos b \cdot \cos c = -\sin^2 A \cdot \sin^2 b \cdot \sin^2 c$$

Using equation 11 yields...

$$\text{Equation 14} \quad 2 - \sin^2 a - \sin^2 b - \sin^2 c - 2 \cdot \cos a \cdot \cos b \cdot \cos c = -\sin^2 B \cdot \sin^2 a \cdot \sin^2 c$$

Using equation 12 yields...

$$\text{Equation 15} \quad 2 - \sin^2 a - \sin^2 b - \sin^2 c - 2 \cdot \cos a \cdot \cos b \cdot \cos c = -\sin^2 C \cdot \sin^2 a \cdot \sin^2 b$$

Equate the right sides of equations 13, 14 and 15 and rearrange...

Law of Sines...

$\frac{\sin a}{\sin A} = \frac{\sin b}{\sin B} = \frac{\sin c}{\sin C}$
---

LAW of COTANGENTS DERIVATION

Equation 16       $\sin b \cdot \sin A = \sin a \cdot \sin B$       ... by Law of Sines

Equation 17       $\cos a = \cos b \cdot \cos c + \sin b \cdot \sin c \cdot \cos A$

Equation 18       $\cos b = \cos a \cdot \cos c + \sin a \cdot \sin c \cdot \cos B$

Equation 19       $\cos c = \cos a \cdot \cos b + \sin a \cdot \sin b \cdot \cos C$

} by Law of Cosines

Substitute for  $\cos b$  from equation 18 into equation 17...

$$\cos a = (\cos a \cdot \cos c + \sin a \cdot \sin c \cdot \cos B) \cdot \cos c + \sin b \cdot \sin c \cdot \cos A$$

$$\cos a = \cos a \cdot \cos^2 c + \sin a \cdot \sin c \cdot \cos c \cdot \cos B + \sin b \cdot \sin c \cdot \cos A$$

$$\cos a - \cos a \cdot \cos^2 c = \sin c \cdot \sin a \cdot \cos c \cdot \cos B + \sin c \cdot \sin b \cdot \cos A$$

$$\cos a \cdot (1 - \cos^2 c) = \sin c \cdot (\sin a \cdot \cos c \cdot \cos B + \sin b \cdot \cos A)$$

Apply the trig identity  $\sin^2 \theta = 1 - \cos^2 \theta$  ...

$$\cos a \cdot \sin^2 c = \sin c \cdot (\sin a \cdot \cos c \cdot \cos B + \sin b \cdot \cos A)$$

$$\cos a \cdot \sin c = \sin a \cdot \cos c \cdot \cos B + \sin b \cdot \cos A$$

Rearrange and divide both sides with equation 16...

$$\frac{\sin b \cdot \cos A}{\sin b \cdot \sin A} = \frac{\cos a \cdot \sin c - \sin a \cdot \cos c \cdot \cos B}{\sin a \cdot \sin B}$$

Law of Cotangents...

$$\cot A = \frac{\cot a \cdot \sin c - \cos c \cdot \cos B}{\sin B}$$

Or ...

$$\tan A = \frac{\sin B}{\cot a \cdot \sin c - \cos c \cdot \cos B}$$

Spherical Excess...

$$\tan\left(\frac{E}{4}\right) = \sqrt{\tan\left(\frac{s}{2}\right) \cdot \tan\left(\frac{s-a}{2}\right) \cdot \tan\left(\frac{s-b}{2}\right) \cdot \tan\left(\frac{s-c}{2}\right)}$$

where

$$s = \frac{a+b+c}{2}$$